arXiv:1611.08739v1 [math.GR] 26 Nov 2016

On just-infinite periodic locally soluble groups

R. Grigorchuk and P. Shumyatsky

Abstract. We construct an uncountable family of periodic locally soluble groups which are hereditary just-infinite. We also show that $C^*(G)$ is just-infinite for many groups G in this family.

1. Introduction

An infinite group G is just-infinite if every proper quotient of G is finite. Just-infinite groups are on the border between finite and infinite groups and therefore deserve special attention. In [**2**] the first author introduced three classes of just infinite groups: branch just infinite groups, and just-infinite groups that contain a subgroup of finite index which is a direct product of finitely many copies of a group L, where L is either simple or hereditary just-infinite. A group L is hereditary just-infinite if it is infinite, residually finite and every subgroup of finite index in L is just-infinite. He showed, using results of Wilson [**5**], that the class of just-infinite groups splits into these three subclasses.

In [**3**] just-infinite C^*-algebras were introduced and studied. In particular, a trichotomy (splitting in three subclasses) was proved, but this trichotomy uses quite different ingredients than in [**2**] (still the simplicity and residual finiteness are involved). In [**3**] a problem about existence of groups with just-infinite group C^*-algebras was discussed. As a possible approach for handling the problem it was suggested to deal first with the question about existence of residually finite, locally finite, hereditary just-infinite groups (see [**3**, Problem 6.10]).

Such a group was constructed in the work of Belyaev and the authors [**1**] thus confirming that locally finite groups with just-infinite

2010 *Mathematics Subject Classification.* 46L05, 20F50, 16S34.

The first author was supported by NSA grant H98230-15-1-0328. The second author was supported by FAPDF and CNPq.

group C^*-algebra exist. On the other hand, results obtain in [**1**] suggest that the structure of a locally finite just-infinite group G must be rather special. Lemma 3.1 in [**1**] implies that such a group G is never locally nilpotent while Theorem 3.3 in [**1**] tells us that G must have infinite exponent. It is an open problem whether G can satisfy a nontrivial law.

In the present article we give first examples of periodic locally soluble groups which are hereditary just-infinite. Actually we construct an uncountable family of such groups and show that $C^*(G)$ is just-infinite for many groups G in this family (see Theorems 2.9 and 2.10 in the next section). Moreover, we note that some of our groups G are rather "close to being a p-group" in the sense that they are $\{p,q\}$-groups, that is, they have the set of prime divisors of orders of their elements containing only two primes.

2. Results

If X and Y are nonempty subsets of a group G, we denote by $[X,Y]$ the subgroup generated by all commutators $[x,y]$, where $x \in X$ and $y \in Y$. Suppose α is an automorphism of G. We write x^α for the image of x under α. If A is a group of automorphisms of G and $H \leq G$, the subgroup generated by elements of the form $h^{-\alpha}h$ with $h \in H$ and $\alpha \in A$ is denoted by $[H,A]$. It is well-known that the subgroup $[G,A]$ is always an A-invariant normal subgroup in G. We also write $C_H(A)$ for the fixed point subgroup of A in H. In the case where A is cyclic generated by α we write $[H,\alpha]$ and $C_G(\alpha)$, respectively. When convenient, we identify elements of the group G with the corresponding inner automorphisms of G. As usual, the symbols G' and $Z(G)$ stand for the derived group and the center of a group G, respectively.

We will require the following well-known fact.

LEMMA 2.1. *Let G be a metabelian group and $a \in G$. Suppose that B is an abelian subgroup in G containing the commutator subgroup G'. Then $[B,a]$ is normal in G.*

PROOF. Since B contains G', it follows that B is normal in G. Let $x \in G$. Write $a^x = ay$ with $y \in G'$. Taking into account that $[B,y] = 1$, we obtain $[B,a]^x = [B,a^x] = [B,ay] = [B,a]$. Thus, an arbitrary element $x \in G$ normalizes $[B,a]$, as required. □

Throughout, we use the following standard facts on coprime groups of automorphisms without explicit references.

LEMMA 2.2. *Let A be a group of automorphisms of the finite group G with $(|A|,|G|) = 1$.*

(1) *If N is any A-invariant normal subgroup of G, we have $C_{G/N}(A) = C_G(A)N/N$;*
(2) *$G = [G, A]C_G(A)$;*
(3) *If G is abelian, then $G = [G, A] \times C_G(A)$;*

LEMMA 2.3. *Let G be a finite group admitting a coprime automorphism α such that $G' \leq C_G(a)$. Set $Z = Z([G, a])$. Then $[Z, a] \leq Z(G)$.*

PROOF. Let $g \in G$ and $h \in C_G(a)$. Taking into account that $[h, g] \in C_G(a)$, write

$$h^g = (h^g)^\alpha = h^{g^\alpha}.$$

We therefore deduce that $[h, [g, \alpha]] = 1$. This shows that $C_G(\alpha)$ commutes with $[G, \alpha]$. Now the lemma follows from the fact that $G = [G, \alpha]C_G(\alpha)$. □

LEMMA 2.4. *Let $G = G_1 \oplus G_2 \oplus \cdots \oplus G_m$ be a finite abelian group written as a direct sum of isomorphic subgroups G_i. Suppose that G admits a coprime automorphism α, of order m, which cyclically permutes the summands G_i. Then the index of $[G, \alpha]$ in G is precisely the order of G_1.*

PROOF. This is immediate from the facts that $G = [G, \alpha] \oplus C_G(\alpha)$ and $C_G(\alpha) = \{x + x^\alpha + \cdots + x^{\alpha^{m-1}}; x \in G_1\}$ is just the diagonal subgroup of G. □

Let p be a prime. Recall that a finite p-group G is extra-special if $G' = Z(G)$ and $G/Z(G)$ is elementary abelian. In the present paper we will work with central products of such groups. We use the term "central product" as in [**4**]: a group G is a central product of its subgroups H and K (written $G = H \circ K$) if G is generated by H and K such that $[H, K] = 1$ and $H \cap K = Z(H) = Z(K)$.

LEMMA 2.5. *Let $G = G_1 \circ G_2 \circ \cdots \circ G_m$ be a finite group which is a central product of isomorphic extra-special subgroups G_i of order p^3. Suppose that G admits a coprime automorphism α, of order m, which cyclically permutes the factors G_i such that $G' \leq C_G(\alpha)$. Then $[G, \alpha]$ is extra-special and has index p^2 in G.*

PROOF. It is clear that G is extra-special. If $[G, \alpha]$ is abelian, then by Lemma 2.3 we have $[G, \alpha] \leq Z(G)$. This leads to a contradiction since $Z(G) = G' \leq C_G(\alpha)$. Hence $[G, \alpha]$ is nonabelian and $[G, \alpha]' = G'$. Further, Lemma 2.3 shows that $Z([G, \alpha]) \leq C_G(\alpha)$. Therefore $[G, \alpha]$ is extra-special.

Let the automorphism of G/G' induced by α be denoted again by the symbol α. Lemma 2.4 shows that the index of $[G/G', \alpha]$ in G/G' is

precisely p^2. Therefore, back in G, the index of $[G,\alpha]G'$ is p^2 and since $G' \leq [G,\alpha]$, we conclude that the index of $[G,\alpha]$ is p^2. □

Given two groups K and M, we write $M \wr K$ for the wreath product of M by K. Thus, $M \wr K$ is a semidirect product of a normal subgroup B, isomorphic to the direct product of $|K|$ copies of M, and the subgroup isomorphic to K. Thus, we will naturally identify K with the subgroup acting on B.

Suppose that M is an extra-special group of order p^3 for some prime p. It is clear that $Z(B) = B'$. Obviously, the subgroup $[B', K]$ is normal in $M \wr K$. Set $B_0 = B/[B', K]$ and let H be the natural semidirect product of B_0 by K. Of course, H is isomorphic to the quotient group $(M \wr K)/[B', K]$. Recall that $B = M_1 \times M_2 \times \cdots \times M_{|K|}$, where the factors M_i are isomorphic with M. Since K transitively permutes the commutator subgroups M_i' and since the subgroup $[B', K]$ is K-invariant, the fact that B_0 is nonabelian implies that no subgroup M_i' is contained in $[B', K]$. Thus, the images of M_i in B_0 are isomorphic with M_i. We therefore from now on will regard M_i as subgroups of B_0. Let $x \in Z(B_0)$. Write $x = x_1 x_2 \dots x_{|K|}$ for suitable $x_i \in M_i$. Suppose that $x_1 \notin B_0'$. Then we can choose $y \in M_1$ such that $[x_1, y] \neq 1$. Then we also have $[x, y] \neq 1$, which contradicts the assumption that $x \in Z(B_0)$. Therefore $x_1 \in B_0'$. Similarly, we deduce that $x_i \in B_0'$ for all i and therefore $Z(B_0) = B_0'$. It follows that B_0 is a central product of the groups M_i. Let $a \in K$ be a p'-element and set $D = [B_0, a]$. In view of Lemma 2.5 we deduce that D is extra-special and $D' = Z(D) = B_0'$.

Let $p_1, p_2, \dots$ be an infinite sequence of primes such that $p_{i+1} \neq p_i$ for all $i = 1, 2, \dots$ (but we allow $p_{i+2} = p_i$). We will construct an infinite sequence of finite soluble groups $G_1, G_2, \dots$ such that for each $i = 1, 2, \dots$ the following holds.

(a) There is a natural embedding $G_i \leq G_{i+1}$.
(b) G_i has a unique minimal normal subgroup R_i;
(c) $R_i \leq Z(G_i)$;
(d) R_i is of order p_i.

Let $G_1 = R_1$ be the cyclic group of order p_1. Now assume that $i \geq 2$ and the group G_{i-1} satisfies the above conditions. Let M_i be an extra-special p_i-group of order p_i^3 and $H = M_i \wr G_{i-1}$. Denote by B the base group of this wreath product. We view G_{i-1} as a subgroup of H. Of course, $[B', H] = [B', G_{i-1}]$. The index of $[B', G_{i-1}]$ in B' is exactly p_i. The group G_{i-1} naturally acts on the quotient group $B_i = B/[B', H]$. Since $R_{i-1} \leq Z(G_{i-1})$, it follows that $D_i = [B_i, R_{i-1}]$ is normalized by G_{i-1}. Let G_i be the natural semidirect product of D_i by G_{i-1}. Since R_{i-1} is the unique minimal normal subgroup in G_{i-1} and since R_{i-1}

does not centralize D_i, it follows that G_{i-1} faithfully acts on D_i and therefore each normal subgroup of G_i has nontrivial intersection with D_i. Taking into account that D_i' is a unique minimal normal subgroup in D_i we conclude that D_i' is a unique minimal normal subgroup in G_i. Therefore we can define $R_i = D_i'$. Thus, the group G_i has the required properties.

For indices $2 \leq k \leq i$ we define the subgroup T_{ik} in G_i by setting $T_{ik} = \prod_{j=k}^{i} D_j$. It is clear that $T_{ii} = D_i$.

LEMMA 2.6. *For a fixed i we have.*

(a) *The subgroups T_{ik} are normal in G_i.*
(b) *Each subgroup T_{ik} is complemented in G_i by the subgroup G_{k-1}.*
(c) $[T_{ik}, R_{k-1}] = T_{ik}$.

PROOF. The lemma is quite obvious when $i = 2$. In the general case all claims can be obtained by an easy induction on i. We therefore omit the details. □

Let G be the union (direct limit) of the above sequence $G_1 \leq G_2 \leq \ldots$.

LEMMA 2.7. *The group G is a locally finite, residually finite, hereditary just-infinite group.*

PROOF. The claim that G is locally finite is obvious. For each $k \geq 2$ we denote by T_k the union of all T_{ik} with $i = k, k+1, \ldots$. Since T_{ik} is normal in G_i, it follows that T_k is normal in G. Moreover, T_k is complemented in G by G_{k-1}. Therefore each T_k has finite index in G. Suppose that x is a nontrivial element lying in T_k for each k. Then x does not belong to G_{k-1} for each k. This is a contradiction since G is the union of the sequence $G_1 \leq G_2 \leq \ldots$. Hence, the intersection of all T_k is trivial and so G is residually finite.

Let N be a proper nontrivial normal subgroup of G. Let j be the minimal index such that $R_j \leq N$. For any $i > j$ the normal closure of R_j in G_i is contained in N. In view of Lemma 2.6(c) we conclude that T_{j+1} is contained in N, as well. Since the index of T_{j+1} in G is finite, so is the index of N. Thus, G is just-infinite and every subgroup of finite index contains the subgroup T_k for some k. Hence, once we show that every normal subgroup of T_k has finite index, we will be able to conclude that G is a hereditary just-infinite group.

Set $T = T_k$ and let S be a normal subgroup of T. Recall that T is the union of the finite subgroups T_{ik} with $i = k, k+1, \ldots$. Let j be the minimal index such that $S \cap T_{jk} \neq 1$. Because of minimality of j we have $S \cap D_j \neq 1$. Taking into account that $R_j = D_j'$ is a unique

normal subgroup of D_j we conclude that R_j is contained in S. Invoking Lemma 2.6(c) deduce that $T_{i(j+1)}$ is contained in S for any $i \geq j+1$. Therefore the subgroup of finite index $T_{(j+1)}$ is contained in S. Hence, S has finite index in G. □

We will now have a closer look at the action of G_{i-1} on D_i. Recall that $D_i = [B_i, R_{i-1}]$, where $B_i = A_1 \circ A_2 \circ \cdots \circ A_{|G_{i-1}|}$ is a central product of extra-special groups A_j of order p_i^3, each isomorphic with M_i, and the group G_{i-1} regularly permutes the factors. Let $O_1, O_2, \ldots, O_m$ be the orbits under the permutational action of R_{i-1} on the set $\{A_1, A_2, \ldots, A_{|G_{i-1}|}\}$. For each $l \leq m$ set

$$E_l = [\prod_{A_j \in O_l} A_j, R_{i-1}].$$

By Lemma 2.5, each E_j is extra-special. Thus,

$$D_i = E_1 \circ E_2 \circ \cdots \circ E_l \qquad (*)$$

is a central product of isomorphic extra-special groups E_j. We see that R_{i-1} normalizes each of the factors E_l and the natural permutational action of G_{i-1}/R_{i-1} on the set $\{E_1, E_2, \ldots, E_l\}$ is a regular one. In what follows this fact will play an important role. We will also need the following observation.

LEMMA 2.8. *Each of the subgroups E_j contains non-central elements of prime order.*

PROOF. Set $E = E_j$ and $R = R_{i-1}$. Thus, by Lemma 2.5, E is an extra-special p_i-group of index p_i^2 in $\prod_{A_j \in O_l} A_j$. Further, E is acted on by the group R, of order p_{i-1}, in such a way that $E = [E, R]$. Suppose that E has no non-central elements of prime order. Then of course, E is isomorphic with the quternion group of order 8 and R has order 3. On the other hand, since R has order 3, the group $\prod_{A_j \in O_l} A_j$ is a central product of 3 extraspecial groups of order 8. Thus $\prod_{A_j \in O_l} A_j$ has order 2^7. Being a subgroup of index 4, the subgroup E must have order 2^5, a contradiction. □

A remarkable property of our groups G is that in many cases all ideals in the group algebra $K[G]$ have finite codimension. More precisely, we have the following theorem.

THEOREM 2.9. *Let K be a field of characteristic p (possibly with $p = 0$). Suppose that the sequence $p_1, p_2, \ldots$ has the property that for any index i there exist indices u, v such that $i < u < v$ and $p_u = p_v \neq p$. Then every ideal in the group algebra $K[G]$ has finite codimension.*

Before we embark on the proof, note that many sequences of primes verify the condition in the theorem. For instance, this happens whenever a prime q, different from p, occures in the sequence infinitely many times, or whenever infinitely many primes occure in the sequence more than once.

Proof of Theorem 2.9. Let I be a nonzero ideal in $K[G]$ and $0 \neq \alpha \in I$. Then we can assume that 1 occurs in the support of α and we write $\alpha = \sum_0^n k_i x_i^{-1}$ with $1 = x_0, x_1, \ldots, x_n$ distinct elements of G and $k_0 \neq 0$. Let t be the minimal index such that $x_0, x_1, \ldots, x_n \in G_t$. Set $X = \langle x_0, x_1, \ldots, x_n \rangle$ and remark that $X \cap R_{t+j} = 1$ for any $j \geq 1$. Choose indices $v > u > t+1$ such that $p_u = p_v = q \neq p$ and consider the decomposition $D_u = E_1 \circ E_2 \circ \cdots \circ E_l$ as in (*). Since $X \cap R_{u-1} = 1$, it follows that X regularly permutes the factors E_i. By Lemma 2.8 E_1 contains a non-central element y of order q. For $i = 0, 1, \ldots, n$ set $y_i = y^{x_i}$ and $z_i = [y_i, x_i]$. Denote the subgroup $\langle y_i^{x_j}; 0 \leq i, j \leq n \rangle$ by A. Since no two of the elements $y_i^{x_j}$ belong to the same factor E_k, it follows that A is abelian and the subgroup $\langle z_1, \ldots, z_n \rangle$ is elementary abelian of order q^n.

We show now by inverse induction on s with $n \geq s \geq 0$ that I contains an element

$$\beta_s = \sum_0^s \beta_{si} x_i^{-1}$$

such that $\beta_{si} \in K[A]$ and, if $s < n$, we have

$$\beta_{s0} = k_0(z_n - 1)(z_{n-1} - 1) \ldots (z_{s+1} - 1).$$

First for $s = n$ we merely take $\beta_s = \alpha$. Now suppose we have β_s as above contained in I with $n > s > 0$. Then $y_s^{-1} \beta_s y_s$ and $z_s \beta_s$ both belong to I and hence

$$\beta_{s-1} = z_s \beta_s - y_s^{-1} \beta_s y_s \in I.$$

Since A is abelian and $y_s, z_s \in A$, we have

$$\beta_{s-1} = \sum_0^s z_s \beta_{si} x_i^{-1} - \sum_0^s y_s^{-1} \beta_{si} x_i^{-1} y_s$$

$$= \sum_0^s \beta_{si}(z_s - [y_s, x_i]) x_i^{-1} = \sum_0^{s-1} \beta_{si}(z_s - [y_s, x_i]) x_i^{-1}$$

since $z_s = [y_s, x_s]$. Recall that $x_0 = 1$. It follows that $\beta_{s-1,0}$ has the required form and the induction step is proved.

In particular, when $s = 0$ we conclude that

$$\beta_0 = k_0(z_n - 1)(z_{n-1} - 1) \ldots (z_1 - 1) \in I.$$

Here $k_0 \neq 0$ and, since $\langle z_1, z_2, \dots, z_n\rangle = \langle z_1\rangle \times \langle z_2\rangle \times \cdots \times \langle z_n\rangle$, we conclude that $\beta_0 \neq 0$. Thus, we have shown that $I \cap K[A] \neq 0$.

Now write $A = \{1 = a_0, a_1, \dots, a_m\}$. We know that $A \cap R_u = 1$ and so $A \cap R_{u+j} = 1$ for any $j \geq 0$. Consider now the decomposition $D_v = \tilde{E}_1 \circ \tilde{E}_2 \circ \cdots \circ \tilde{E}_{l'}$ as in (*). The group A regularly permutes the factors $\tilde{E}_j$. In view of Lemma 2.8 we can choose a non-central element b of order q that belongs to $\tilde{E}_1$. For $i = 0, 1, \dots, m$ set $b_i = b^{a_i}$. Denote the subgroup $\langle b_0, b_1, \dots, b_m\rangle$ by B. Set $H = BA$. The choice of the element b guarantees that $H \cong C_q \wr A$, where C_q denotes the cyclic group of order q. In particular, the center $Z(H)$ is cyclic generated by the product $b_0 \cdots b_m$.

Fix $i \leq m$ and set $B_i = [B, a_i]$. Since B is abelian, B_i is isomorphic with $B/C_B(a_i)$. Let $\hat{B}_i$ denote the sum of elements in B_i. We compute

$$\sum_{b\in B}(a_i^{-1})^b = \sum_{b\in B}[b, a_i]a_i^{-1} = |C_B(a_i)|\hat{B}_i a_i^{-1}.$$

Let $\alpha_0 = \sum l_i a_i^{-1} \in I \cap K[A]$ with $l_0 \neq 0$. Then $\beta = \sum_{b\in B} b^{-1}\alpha_0 b \in I$. By the above

$$\beta = \sum_i l_i |C_B(a_i)|\hat{B}_i a_i^{-1}.$$

For $i = 0$, since $a_0 = 1$, the summand here is $l_0|B|$. For $i \neq 0$ the element a_i does not centralize B so $B_i \neq 1$. Lemma 2.1 tells us that each subgroup B_i is normal in H. Recall that $Z(H)$ is cyclic of order q. Because H is nilpotent, every normal subgroup of H has nontrivial intersection with $Z(H)$. Consequently, $Z(H) \leq B_i$. Let z be a generator of $Z(H)$. Since $z \in B_i$, we have $(z-1)\hat{B}_i = 0$. Therefore

$$(z-1)\beta = l_0|B|(z-1).$$

We put this together with the facts that $(z-1)\beta \in I$ and $l_0 \neq 0$. Since the characteristic of K is not q, it follows that $z - 1 \in I$.

Now set $N = \{g \in G; g - 1 \in I\}$. This is a normal subgroup in G. The above shows that $N \neq 1$. Since G is just-infinite, we conclude that N has finite index. This implies that I has finite codimension. □

Combining Theorem 2.9 with results from [**3**] we can now easily show that whenever G is as in Theorem 2.9 with $p = 0$ the C^*-algebra $C^*(G)$ is just-infinite. We denote by $\mathbb{C}$ the field of complex numbers.

THEOREM 2.10. *Assume that the sequence $p_1, p_2, \dots$ has the property that for any index i there exist indices u, v such that $i < u < v$ and $p_u = p_v$. Then $C^*(G)$ is just-infinite.*

PROOF. By Theorem 2.9, the group algebra $\mathbb{C}[G]$ is just-infinite. The result now follows immediately from Propositions 6.2 and 6.6 of [**3**]. □

References

[1] V. Belyaev, R. Grigorchuk, P. Shumyatsky, On just-infiniteness of locally finite groups and their C^*-algebras, Bull. Math. Sci., to appear, doi:10.1007/s13373-016-0091-4.

[2] R. I. Grigorchuk, Just infinite branch groups. New horizons in pro-p groups, 121–179, Progr. Math., **184**, Birkhäuser Boston, Boston, MA, 2000.

[3] R. Grigorchuk, M. Musat, M. Rørdam, Just-infinite C^*-algebras, Preprint, 2016, https://arxiv.org/abs/1604.08774.

[4] C. R. Leedham-Green, S. McKay, The structure of groups of prime power order, London Mathematical Society Monographs. New Series, 27 (2002), Oxford University Press.

[5] J. S. Wilson, Groups with every proper quotient finite, Proc. Cambridge Philos. Soc., **69** (1971), 373–391.

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX, USA

E-mail address: grigorch@math.tamu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRASÍLIA, 70910 BRASÍLIA DF, BRAZIL

E-mail address: pavel@unb.br